

Publications: full papers

- 1) The action of Acetylcholine and L-glutamic acid on rat caudate neurones.  
G. Bernardi, V. Floris, M.G. Marciani, C. Morocutti and P. Stanzione.  
Brain Res. 1976 (114), 134-138.
- 2) The action of Dopamine on rat caudate neurones intracellularly recorded.  
G. Bernardi, M.G. Marciani, C. Morocutti, F. Pavone and P. Stanzione.  
Neurosciences Letters 1978 (8), 235-240.
- 3) The inhibitory action of Glycine on rat cortical neurones.  
G. Bernardi, E. Cherubini, M.G. Marciani, P. Stanzione.  
Neuroscience Letters 1979 (12), 335-338.
- 4) Action mechanism of gamma aminobutyric acid (GABA) and Glycine on rat cortical neurones.  
M.G. Marciani, P. Stanzione, E. Cherubini, G. Bernardi.  
Neuroscience Letters 1980 (18), 169-172.
- 5) Evidence in favour of Gaba as an inhibitory transmitter in rat corpus striatum.  
G. Bernardi, M.G. Marciani, P. Stanzione, E. Cherubini, N. Mercuri.  
In " Gaba and the basal ganglia" eds. G. Di Chiara and G.L. Gessa , Raven Press New York 1981 - 69-77.
- 6) The action of Glycine on rat epileptic foci.  
E. Cherubini, P. Stanzione, M.G. Marciani, N. Mercuri.  
Neuroscience Letters 1981 (21), 93-97.
- 7) Evidence in favor of a neurotransmitter role of Glycine in the rat cerebral cortex.  
G. Levi, G. Bernardi, E. Cherubini, V. Gallo, M.G. Marciani, P. Stanzione.  
Brain Res. 1982 (236), 267-131.

8) Responses of intracellularly recorded cortical neurones to the iontophoretic application of dopamine.

G. Bernardi, E. Cherubini, M.G. Marciani, N. Mercuri and P. Stanzione.

Brain Res. 1982 245(2), 267-274.

9) Action of Neurotensin on rat spinal cord neurones.

P. Stanzione, W. Ziegelgansberger.

Brain Res. 1983 (268), 111-118

10) Effect of Dopamine on the threshold of the voltage dependent ionic-channels in the rat brain.

G. Bernardi, P. Calabresi, N. Mercuri and Stanzione P.

Annali Istituto Superiore di Sanità - Roma.

1984 20, ( 1), 1-4.

11) Dopamine decreases the amplitude of excitatory post-synaptic potentials in rat striatal neurons.

Bernardi G., Calabresi P., Mercuri N. and Stanzione P.

In "Basal ganglia structure and function", Eds. J.S. McKenzie, R.E. Kemm, L.N. Wilcock: Plenum Publishing Corporation. New York, 1984, 161-172.

12) Dopamine modulates Cal hippocampal neurons by elevating the threshold for spikes generation: an "in vitro" study.

Stanzione P., Calabresi P., Mercuri N. and Bernardi G.

Neuroscience 1984 13, ( 4), 1105-1116.

13) Dopaminergic and noradrenergic responses in the hippocampal slice preparation. Evidence for different receptors.

Marciani M.G., Calabresi P., P. Stanzione and Bernardi G.

Neuropharmacology 1984 23, ( 3), 303-307.

14) Effectiveness of Bornaprine on parkinson tremor.

Sancesario G., Cicardi M.C., Fiermonte G., Giacomini P. and Stanzione P.

Ital. J. Neurol. Sci. 1984 , V, 289-293.

15) Electrical stimulation of mesencephalic cell groups (A9-A10) produces monosynaptic excitatory potentials in rat frontal cortex.

Mercuri N., Calabresi P., Stanzione P. and Bernardi G.

Brain Res. 1985, (338), 192-195.

16) Dopamine decreases cell excitability in rat striatal neuron by pre and post synaptic mechanisms.

Mercuri N., Bernardi G., Calabresi P., Cotugno A., Levi G. and Stanzione P.

Brain Res.1985 ( 358), 110-121.

17) The excitatory action of morphine on rat cortical neurons is mediated by a postsynaptic mechanism.

Mercuri N., Calabresi P., Stanzione P., Valenti G. and Bernardi G.

Regulatory Peptides 1985, (1), 197-199.

18) Nervous impulse propagation along peripheral and central (spinal and intracranial) fast conducting motor and sensory pathways in man.

Rossini P., Di Stefano E. and Stanzione P.

Electroenceph. clin. Neurophysiol. 1985 (60), 320-334.

19) Intracellular study on the action of dopamine in the mammalian central nervous system. A survey.

Bernardi G., Calabresi P., Mercuri N. and Stanzione P.

In "Modulation of central and peripheral transmitter function". Symposia in Neuroscience. Eds. Biggio G., Spano P.F., Toffano G. and Gessa G.L. Liviana Press - Springer Verlag 1986, (3), 75-80.

20) Action of morphine on rat cortical neurons recorded in vivo: evidence for an excitatory post synaptic effect, naloxone insensitive.

Bernardi G., Calabresi P., Mercuri N. and Stanzione P.

Neuroscience 1986, (3), 126-138.

21) Neurotransmitter mechanisms and cognitive function.

P. Stanzione, P. Calabresi, N. Mercuri, A. Stefani, G. Bernardi.

In "Maturation of the CNS and evoked potentials" Excerpta Medica 1986, 230-235.

22) Intracellular studies on the dopamine induced firing inhibition of neostriatal neurons in vitro: evidence for D1 receptors involvement.

P. Calabresi, N. Mercuri, P. Stanzione, A. Stefani and G. Bernardi.

Neuroscience 1987, 20, (4), 757-771.

23) Morphine induces bursting activity in rat cortical neurons: an intracellular study in vivo.

P. Stanzione, P. Calabresi, N. Mercuri and G. Bernardi.

In "Inactivation of hypersensitive neurons" - N. Chalazonitis and M. Gola Eds. Alan R. Liss - New York 1987, 57-61.

24) Role of D1 and D2 dopamine receptors in the mammalian striatum: electrophysiological studies and functional implications.

Calabresi P., Mercuri N., Stanzione P. and Bernardi G.

In "The Basal Ganglia II" - M.B. Carpenter and A. Jayaraman Eds. Plenum Publishing Corp. 1987

25) Electrophysiological (PERG, VEP) abnormalities in Parkinson disease are reversed by L-Dopa.

Pierelli F., Stanzione P., Peppe A., Stefano E., Rizzo P.A., Bernardi G., Morocutti C.

In "Dopaminergic mechanism in vision". I. Bodis-Wollner and M. Piccolino Eds. Alan R. Liss Inc. New York, 1988, 253-265.

26) Excitatory amino acids in synaptic excitation of rat striatal neurons in vitro.

E. Cherubini, P.L. Herrling, L. Lanfumey and P. Stanzione.

J. Physiol. London 1988, (400), 677-690.

27) Morphine induces a spontaneous and evoked bursting activity in rat cortical neurones by adding a postsynaptic active mechanism to the synaptic input: an intracellular study in vivo.

Stanzione P., Stefani A., Bernardi G.

Neuroscience 1988, 26, (1), 45-53.

28) Met- and Leu-enkephalins inhibit rat cortical neurons intracellularly recorded in vivo while morphine excites them. Evidences for naloxone-sensitive and naloxone-insensitive effects.

Stanzione P., Stefani A., Calabresi P., Mercuri N. and Bernardi G.

Exp. Brain Res. 1989, (77), 302-308.

29) Pattern visual evoked potentials and electroretinogram abnormalities in Parkinson's disease: effects of L-DOPA therapy.

Stanzione P., Pierelli F., Peppe A., Stefano E., Rizzo P.A., Morocutti.

Clin. Vis. Sci. 1989, (4), 115-127.

30) Abnormalities of short latency somatosensory evoked potentials in parkinsonian patients.

P.M. Rossini, F. Babiloni, G. Bernardi, L. Cecchi, P.B. Johnson, A. Malentacca, P. Stanzione and A. Urbano.

Electroenceph. clin. Neurophysiol. 1989, (74), 277-289.

31) Infarti cerebrali di confine in un caso di trombocitemia essenziale.

M. Silvestrini, M. Tagliati, R. Floris, P. Stanzione.

Rivista di Neurologia, 1989, (59), 220-222.

32) Epileptic activity following 10 minute cerebral ischemia in Mongolian gerbils: an electropysiological study.

M.G. Marciani, G. Santone, G. Sancesario, R. Massa, P. Stanzione and G. Bernardi.

Neuroscience Letters 1990, (112), 48-53.

33) Spontaneous subarachnoid hemorrhage in a HIV patient. Silvestrini M., Floris R., Tagliati M., Stanzione P., Sancesario G.

Ital. J. Neurol. Sci. 1990, (11), 493-495.

34) Retinal visual dysfunction, in Parkinson's disease, is enhanced by low contrast stimuli.

P. Stanzione, M. Tagliati, A. Peppe, G. Sancesario, F. Pierelli

In: Basal Ganglia III. G. Bernardi, M. Carpenter, G. Di Chiara, P. Stanzione

(Eds.) Plenum Press, New York, 1991, 630-638.

35) Dopaminergic pharmacological manipulations in normal humans confirm the specificity of the visual ( PERG-VEP ) and cognitive ( P300) electrophysiological alterations in Parkinson's disease.

P. Stanzione, F. Fattapposta, M. Tagliati, C. D'Alessio, M.G. Marciani, A. Foti and G. Amabile.

In: New Trends and Advanced Techniques in Clinical Neurophysiology. F. Mauguiere and P. M.

Rossini Eds. Elsevier, Amsterdam, 1990, 216-220

36) Haloperidol delays pattern electroretinograms more than VEPs in normal humans: a pharmacological tool in Parkinson's disease pathophysiology.

P. Stanzione, M. Tagliati, M. Silvestrini, S. Porcu and G. Bernardi

Clin: Vis. Sci. 1990, ( 6), 137-147.

37) Basal Ganglia III - G. Bernardi, Carpenter M., Di Chiara G., Morelli M. and Stanzione P.

Plenum Press, New York London 1991

38) Epileptic activity following cerebral ischemia in mongolian gerbils is depressed by CPP, a competitive antagonist of the N-methyl-D-aspartate receptor.

M.G. Marciani, Santone G., Sancesario G., Massa R., Stanzione P. and Bernardi G.

Neuroscience Letters 1991 (129) 306-310

39) P300 variations in Parkinsonian patients before and during dopaminergic monotherapy: a suggested dopamine component in P300.

Stanzione P., Fattapposta F., Giunti P., D'Alessio C., Tagliati M., Affricano C. and Amabile G.

Electroencephalography and Clinical Neurophysiology, 1991 (80) 446-453

40) Low contrast stimuli enhance PERG sensitivity to the visual dysfunction in Parkinson's disease.

Peppe A., Stanzione P., Pierelli F., Stefano E., Rizzo P.A., Tagliati M. and Morocutti C.

Electroenceph. clin. Neurophysiol: 1992, ( 822 ), 453-57

41) An electrophysiological study of D2 dopaminergic actions in normal human retina: a tool in Parkinson's disease.

Stanzione P., Traversa R., Pierantozzi M., Semprini R., Marciani M.G. and Bernardi G.

Neuroscience Letters 1992, (136 ), 125-128.

42) Leukocyte aggregation: a possible link between infection and ischemic stroke.

Galante A., Pietroiusti A., Silvestrini M., Stanzione P. and Bernardi G. - Stroke 1992, (23), 1533.

43) Leukocyte aggregation in acute cerebrovascular disease.

Galante A., Silvestrini M., Stanzione P., Pietroiusti A., Baldoni F., Domenici B. and Bernardi G.

Acta Neurol. Scand. 1992, ( 86 ), 446-449.

44) Serial determinations of leukocyte aggregation in patients with ischemic stroke.

Galante A., Pietroiusti A., Magrini A., Floris R., Simonetti G., Silvestrini M., Stanzione P., Bernardi G.

Intern J. Neuroscience 1993, ( 73 ), 69-76.

45) Protective effect of clonazepam on ischemic brain damage induced by 10 minute bilateral carotid occlusion in mongolian gerbils.

Marciani M.G., Santone G., Sancesario G., Massa R., Stanzione P., Bernardi G.

Functional Neurology, 1993, (23), 185-191

46) Parkinson's Disease and Somatosensory Evoked Potentials: Apomorphine induced transient potentiation of frontal components.

Rossini P.M., Traversa R., Boccasena P., Martino G., Passarelli S., Pacifici L., Bernardi G. and Stanzione P.

Neurology, 1993 (43), 2495-2500

47) Spatial frequency tuning of the monkey pattern ERG depends on D2 receptor-linked action of dopamine.

Tagliati M., Bodis-Wollner I., Kovanecz I., Stanzione P.

Vision Research 1994, (34), 2051-2057.

48) PERG modifications in post traumatic patients with extrapyramidal signs and in Parkinson's disease patients.

Peppe A., Stanzione P., Formisano R., Santilli V. and Bernardi G.

In: Dementia in Parkinson's disease. Korczyn A.D. ed. Monduzzi Press-Italy 1994, pp 59-65.

49) P3 components evoked by unilateral or bilateral somatosensory stimulation: normative data for the study of neglect syndrome.

Stanzione P., Santilli A.M., Caltagirone C., Zarola F., Pierantozzi M. and Bernardi G.

Functional Neurology 1994, (6), 145-152.

50) N30 Somatosensory Evoked Potentials in Parkinson's disease: a pharmacological study. .

Traversa R., Pierantozzi M., Semprini R., Loberti M., Cicardi M.C., Stanzione P.

In: Old and new dopamine agonists in Parkinson's disease. Bonuccelli and Rabey eds.

Journal of Neural Transmission. 1995 Suppl. (45),177-185.

51) Increasing doses of l-sulpiride reveal a dose and a spatial frequency-dependent effect of D2 selective blockade in the human electroretinogram.

Stanzione P., Pierantozzi M., Semprini R., Tagliati M., Traversa R. and Bernardi G.

Vision Research. 1995, (35, 18), 2659-2664.

52) PERGs recorded in "de novo" Parkinson's disease patients are more delayed and reversed by L- Dopa therapy than VEPs.

Peppe A., Stanzione P., Pierelli F., De Angelis D., Pierantozzi M. and Bernardi G.

Neurology 1995, (48), 2495-2500.

53) Multimodal evoked potentials in HIV-seropositive patients: relationship between the immune impairment and the neurophysiological function.



Pierelli F., C. Carruba, G. Tilia, L. Parisi, F. Fattaposta, G. Pozzessere, G. Soldati, P. Stanzione, G. D'offizi, I. Mezzaroma.

Acta Neurologica Scandinavica. 1996, (93,4), 266-271.

54) N30 wave of somatosensory evoked potentials in Parkinson's Disease: a pharmacological approach.

Traversa R., Pierantozzi M., Semprini M., Loberti M., Peppe A., Cicardi M.C., and Stanzione P.

In Functional Neuroscience. C. Barber, G. Celesia, G.C. Comi and F. Mauguiere Eds. Elsevier Science S.V. Amsterdam 1996 pg 290-297.

55) N30 wave of somatosensory evoked potentials in Parkinson's Disease: a pharmacological approach.

Traversa R., Pierantozzi M., Semprini M., Loberti M., Peppe A., Cicardi M.C., and Stanzione P.

EEG J. Supp 1996, (46) 193-200.

56) Quantitative EEG changes in non demented Parkinson disease patients before and during l-dopa therapy.

Stanzione P., Marciani M.G., Maschio M., Bassetti M.A., Spanedda F., Pierantozzi M., Semprini R and Bernardi G.

European Journal of Neurology. 1996, (3) 354-362.

57) SEPs N30 amplitude in Parkinson's disease and in pharmacologically induced rigidity: relationship with the clinical status.

P. Stanzione, R. Traversa, M. Pierantozzi, R. Semprini, M. Loberti, A. Peppe, A. Santilli and G. Bernardi.

European J of Neurology 1997, (4) 1-15.

58) EEG changes induced by vigabatrin monotherapy in focal epilepsy.

M.G. Marciani, P. Stanzione, M. Maschio, F. Spanedda, M.A. Bassetti, D. Mattia, G. Bernardi.

Acta Neurol Scand. 1997, (92) 115-120.

59) Effects of increasing doses of apomorphine during stereotaxic neurosurgery in Parkinson's disease: clinical score and internal Globus Pallidus activity.

Stefani, A., Stanzione, P., Bassi, A., Mazzone, P., Vangelista, T., Bernardi, G.

J. Neural Transm. 1997; 104(8/9): 895-904.

60) Age and stage dependency of P300 latency alterations in non demented Parkinson's disease patients without therapy.

Stanzione, P., Semprini R., Pierantozzi M., Santilli A.M., Fadda L., Traversa R., Peppe A. and Bernardi G.

Electroenceph. clin. Neurophysiol. 1998, (108), 80-91.

62) Lamotrigine add-on therapy in focal epilepsy: electroencephalographic and neuropsychological evaluation.

Marciani M.G., Stanzione P., Mattia D., Spanedda F., Bassetti M.A., Maschio M. and Bernardi G.

Clin. Neuropharmacol. 1998; 21(1): 41-47.

63) D1 agonist CY209-243 attenuates the pattern electroretinogram to low spatial frequency stimuli in the monkey.

Peppe A., Antal A., Tagliati M., Stanzione P., Bodis-Wollner I

Neurosci. Lett 1998 (243) 5-8.

64) Does PERG spatial tuning alteration in Parkinson's disease depend on motor disturbances or retinal dopaminergic loss ?

Peppe, A., Stanzione, P., Pierantozzi, M., Semprini R., Bassi A., Santilli A.M., Formisano R., Piccolino M. and Bernardi G.

Electroenceph. clin. Neurophysiol. 1998 (106), 374-382.

65) Antiparkinsonia and anti-LID effects can be obtained by stimulating the same site within the GPi in Parkinson's disease.

Stanzione P., Mazzone P., Peppe A., Pierantozzi M., Stefani A., Bassi A. and Bernardi G.

Neurology 1998 (51), 1776-1777.

66) Electrophysiological and clinical desensitization to apomorphine administration in parkinsonian patients undergoing stereotaxic neurosurgery.

Stefani A., Mazzone P., Bassi A., Bernardi G., Altibrandi M.G., Peppe A., Pierantozzi M. and Stanzione P.

Exp. Neurol. 1999; 156(1); 209-213.

67) A mixed D1 and D2 antagonist does not replay pattern electroretinogram alterations observed with a selective D2 antagonist in normal humans: relationship with Parkinson's disease pattern electroretinogram alterations.

Stanzione P., Bodis-Wollner I., Pierantozzi M., Pierantozzi M., Semprini R., Tagliati M., Peppe A and Bernardi G.

Electroenceph. clin. Neurophysiol. 1999; 110: 82-85.

68) Basal ganglia and gait control: apomorphine administration and internal pallidum stimulation in Parkinson's disease.

Grasso R, Peppe A, Stratta F, Angelini D, Zago M, Stanzione P, Laquaniti F.

Exp. Brain Res. 1999; 126: 139-148.

69) The effects of deep brain stimulation on the frontal N30 component of somatosensory evoked potentials in advanced Parkinson's disease patients.

Pierantozzi M., Mazzone P., Bassi A., Rossini P.M., Peppe A., Altibrandi M.G., Stefani A., Stanzione P.

Electroenceph. clin. Neurophysiol. 1999; 110: 1-8.

70) Contemporary Bilateral targeting of GPi and STN by "Maranello" stereotactic system for deep brain stimulation in Parkinson's disease.

Mazzone P, Stanzione P, Stefania A, Bassi A, Gattoni G, Altibrandi MG, Peppe A, Pierantozzi M, Vangelista S. Bernardi G.

11<sup>th</sup> European Congress of Neurosurgery. Monduzzi Editore 729-734.

71) Pierantozzi M, Sabato A, Leonardis F, Marciani MG, Giacomini P, Bernardi G and Stanzione P. Curariform peripheral block of muscular tone selectively increases pre-central N30 SEPs

component. a pharmacological study carried out on healthy subjects and parkinsonian syndromes.

Exp Brain Res 2000; 133: 368-376.

**IF. 2.306**

72) Fedele E., Mazzone P., Stefani A., Pierantozzi M., Giacomini P., Raiteri M., Bernardi G. and Stanzione P. Microdialysis in Parkinson's disease patients basal ganglia: apomorphine-induced clinical and electrophysiological effects not paralleled by changes in the release of neuroactive amino acids.

Exp. Neurol. 2001, 167: 356-365.

**IF. 3.503**

73) Fedele E., Stefani A., Bassi A., Pepicelli O., Raiteri M., Altibrandi MG., Frasca S., Pierantozzi M., Giacomini P., Stanzione P. and Mazzone P. Apomorphine-induced clinical and electrophysiological effects in Parkinson's disease patients, not paralleled by amino acids release changes: a microdialysis study.

Funct. Neurol. 2001; 16: 57-66.

**IF. 0.463**

74) Brusa L., Pierantozzi M., Peppe A., Altibrandi MG., Giacomini P., Mazzone P. and Stanzione P. Deep brain stimulation (DBS) attentional effects parallel those of l-dopa treatment.

J. Neural Transm. 2001; 108: 1021-1027.

**IF. 2.93**

75) M. Pierantozzi, G. Sancesario, M.G. Marciani, S. Frasca, G. Lunardi, E., Fedele, P. Giacomini, A. Pietroiusti, A. Galante and P. Stanzione. ) Reduced l-dopa absorption and increased clinical fluctuations in Helicobacter pylori-infected Parkinson's disease patients.

Neurological Science 2001; 22: 89-91.

**IF. 0.38**

76) Pierantozzi M, Pietroiusti A, Galante A, Sancesario G, Lunardi G, Fedele E, Giacomini P, Stanzione P. Helicobacter Pylori-induced reduction of acute levodopa absorption in Parkinson's disease patients.

Ann Neurol 2001; 50 (5): 686-687.

**IF. 8.481**

77) Pierantozzi M, Palmieri MG, Marciani MG, Bernardi G, Giacomini P, Stanzione P. Effect of apomorphine on cortical inhibition in Parkinson's disease patients: a transcranial magnetic stimulation study.

Exp Brain Res 2001; 141: 52-62.

**IF. 2.306**

78) Peppe A, Pierantozzi M, Altibrandi MG, Giacomi P, Stefani A, Bassi A, Mazzone P, Bernardi G, Stanzione P. Bilateral GPi DBS is useful to reduce abnormal involuntary movements in advanced Parkinson's disease patients, but its action is related to modality and site of stimulation.

Eur J Neurol 2001; 8: 579-586.

**IF. 2.437**

79) Stefani A., Bassi A., Mazzone P., Pierantozzi M., Gattoni G., Altibrandi MG., Giacomini P., Peppe A., Bernardi G., Stanzione P. Subdyskinetic apomorphine responses in globus pallidus and subthalamus of parkinsonian patients: lack of clear evidence for the "indirect pathway".

Clin Neurophysiol. 2002 Jan;113(1):91-100.

**IF. 2.718**

80) Pierantozzi M, Palmieri MG, Mazzone P, Marciani MG, Rossini PM, Stefani A, Peppe A, Giacomini P, Stanzione P. Deep brain stimulation of both subthalamic nucleus and internal globus pallidus restores intracortical inhibition in Parkinson's disease paralleling apomorphine effects: a paired magnetic stimulation study.

Clin Neurophysiol. 2002 Jan;113(1):108-13.

**IF. 2.718**

81) A. Bassi, E. Fedele, A. Peppe, M. Pierantozzi, G. Gattoni, G. Bernardi, P. Stanzione and A. Stefani. ) Electrophysiological and biochemical changes related to high frequency deep brain stimulation in Parkinson's disease

In Basal Ganglia Structure and Function VIII - IBAGS VIII- Richard Faull and Luoise Nicholson eds. Pergamon Press 2002

82) E. Troisi, A. Peppe, M. Pierantozzi., M. Matteis, F. Vernieri, P. Stanzione, M. Silvestrini, C. Caltagirone. Emotional processing in Parkinson's disease. A study using functional transcranial doppler sonography.

J. Neurol. 2002; 249: 993-1000.

**IF. 2.653**

- 83) Brusa L, Bassi A, Pierantozzi M, Gaudiello S, Frasca F, Floris R, Stanzione P. Perfusion-weighted dynamic susceptibility (DSC) MRI: basal ganglia hemodynamic changes after apomorphine in Parkinson's disease.  
Neurol Sci. 2002; 23 Suppl 2:S61-2. **IF. 0.38**
- 84) Finazzi-Agro E, Peppe A, D'Amico A, Petta F, Mazzone P, Stanzione P, Micali F, Caltagirone C. Effects of subthalamic nucleus stimulation on urodynamic findings in patients with Parkinson's disease.  
J Urol. 2003; 169: 1388-91. **IF. 3.19**
- 85) Brusa L, Bassi A, Stefani A, Pierantozzi M, Peppe A, Caramia MD, Boffa L, Ruggieri S, Stanzione P. Pramipexole in comparison to l-dopa: a neuropsychological study.  
J Neural Transm. 2003; 110: 373-80. **IF. 2.932**
- 86) di Michele F, Longone P, Romeo E, Lucchetti S, Brusa L, Pierantozzi M, Bassi A, Bernardi G, Stanzione P. Decreased plasma and cerebrospinal fluid content of neuroactive steroids in Parkinson's disease.  
Neurol Sci. 2003 Oct;24(3):172-3. **IF. 0.38**
- 87) Formisano R, Voogt RD, Buzzi MG, Vinicola V, Penta F, Peppe A, Stanzione P. Time interval of oral feeding recovery as a prognostic factor in severe traumatic brain injury.  
Brain Inj. 2004 Jan;18(1):103-9. **IF. 1.182**
- 88) Koch G., Brusa L., Caltagirone C., Olivieri M., Peppe A., Tiraboschi P., and Stanzione P. Subthalamic deep brain stimulation improves time perception in Parkinson's disease.  
Neuroreport 2004; 15(6): 1071-73. **IF. 2.374**
- 89) Antonella Peppe, Mariangela Pierantozzi, Andrea Bassi, Maria Grazia Altibrandi, Livia Brusa, Alessandro Stefani, Paolo Stanzione and Paolo Mazzone. Stimulation of the Subthalamic Nucleus compared to the Globus Pallidus Internus In Parkinson's Disease Patients.  
J Neurosurg. 2004; 101(2): 195-200. **IF. 2.748**

90) Pireantozzi M., Marciani MG., Palmieri MG., Brusa L., Galti S., Caramia MD., Bernardi G., Stanzione P. Effect of vigabatrin on motor responses to transcranial magnetic stimulation. An effective tool to investigate in vivo GABAergic cortical inhibition in humans.

Brain Res 2004; 1028: 1-8.

**IF.2.489**

91) Pierantozzi M., Panella M., Calmieri MG., Koch G., Giordano A., Marciani MG., Bernardi G., Stanzione P., Stefani A. Different TMS pattern of intracortical inhibition in early onset Alzheimer dementia and frontotemporal dementia.

Clinical Neurophysiology 2004; 115(10): 2410-18.

**IF. 2.718**

92) Brusa L., Bassi A., Lunardi GL, Fedele E., Peppe A., Stefani A., Pasqualetti P., Stanzione P., Pierantozzi M. Delayed administration may improve entacapone effects in parkinsonian patients non-responding to the drug.

Eur. J Neurol. 2004; 11(9): 593-606.

**IF. 2.437**

93) Brusa L., Tiraboschi P., Koch G., Peppe A., Pierantozzi M., Ruggeri S., Stanzione P. Pergolide effect on cognitive function in early-mild Parkinson's disease.

J. Neural Transm. 2004; 10:

**IF. 2.932**

94) Forte G., Bocca B., Senofonte O., Petrucci F., Brusa L., Stanzione P., Zannino S., Violante N., Alimenti A., Sancesario G. Trace and major elements in whole blood, serum, cerebrospinal fluid and urine of patients with Parkinson's disease.

J. Neural Transm. 2004; 111(8): 1031-40.

**IF. 2.932**

95) Brusa L., Pierantozzi M., Bassi A., Fedele E., Lunardi G., Giacomini P., Stanzione P. Temporale administration of entacapone with slow release l-dopa: pharmacokinetic profile and clinical outcome.

Neurol. Sci 2004; 25(2): 53-56.

**IF. 0.38**

96) Romigi A., Brusa L., Marciani MG, Pierantozzi M., Placidi F., Izzi F., Sperli F., Testa F., Stanzione P. Sleep episodes and day-time somnolence as result of individual susceptibility to different dopaminergic drugs in a PD patient: a polysomnographic study.

J Neurol Sci. 2005 Jan 15;228(1):7-10.

**I.F. 2.324**

97) Centonze D., Palmieri MP., Boffa L., Pierantozzi M., Stanzione P., Brusa L., Marciani MG., Siracusano A., Bernardi G., Caramia MD. Cortical hyperexcitability in posttraumatic stress disorder secondary to accidental head trauma: a neurophysiological study.

J Psychiatry Neurosci. 2005 Mar;30(2):127-32.

**I.F. 3.579**

98) Ghezzi D, Marelli C, Achilli A, Goldwurm S, Pezzoli G, Barone P, Pellecchia MT, Stanzione P, Brusa L, Bentivoglio AR, Bonuccelli U, Petrozzi L, Abbruzzese G, Marchese R, Cortelli P, Grimaldi D, Martinelli P, Ferrarese C, Garavaglia B, Sangiorgi S, Carelli V, Torroni A, Albanese A, Zeviani M. Mitochondrial DNA haplogroup K is associated with a lower risk of Parkinson's disease in Italians.

Eur J Hum Genet. 2005; 13(6): 748-52.

**IF. 3.564**

99) Koch G., Brusa L., Olivieri M., Stanzione P., Caltagirone C. Memory for time intervals is impaired in left hemi-Parkinson patients.

Neuropsychologia. 2005;43(8):1163-7.

**IF. 4.345**

100) G. Koch, M. Oliveri, L. Brusa, P. Stanzione, S. Torriero, and C. Caltagirone. High frequency rTMS improves time perception in Parkinsons disease.

Neurology. 2005 Aug 23;65(4):623-5.

**IF. 8.172**

101) A. Stefani, E. Fedele, S. Galati, O. Pepicelli, S. Frasca, M. Pierantozzi, A. Peppe, L. Brusa, A. Orlacchio, A.H. Hainsworth, G. Gattoni, P. Stanzione, G. Bernardi, M. Raiteri, and P. Mazzone. Subthalamic Stimulation Activates Internal Pallidus: Evidence from cGMP Microdialysis in PD Patients.

Ann Neurol 2005;57:448-452

**IF. 9.317**

102) Schillaci O, Pierantozzi M, Filippi L, Manni C, Brusa L, Danieli R, Bernardi G, Simonetti G, Stanzione P. The effect of levodopa therapy on dopamine transporter SPECT imaging with(123)I-FP-CIT in patients with Parkinson's disease.

Eur J Nucl Med Mol Imaging. 2005 Sep 8;

**IF. 4.531**



103) Pisani A, Fezza F, Galati S, Battista N, Napolitano S, Finazzi-Agro A, Bernardi G, Brusa L, Pierantozzi M, Stanzione P, Maccarrone M. High endogenous cannabinoid levels in the cerebrospinal fluid of untreated Parkinson's disease patients.

Ann Neurol. 2005 May;57(5):777-9.

**IF. 9.317**

104) G. Koch, L. Brusa, C. Caltagirone, A. Peppe, M. Oliveri P. Stanzione, D Centonze. RTMS of supplementary motor area modulates therapy-induced dyskinesias in Parkinson's disease.

Neurology 2005; 65(4): 623-5.

**IF. 8.172**

105) Brusa L, Versace V, Koch G, Bernardi G, Iani C, Stanzione P, Centonze D. Improvement of choreic movements by 1 Hz repetitive transcranial magnetic stimulation in Huntington's disease patients.

Ann Neurol. 2005; 58(4); 655-6.

**IF. 9.317**

106) Mazzone P, Lozano A, Stanzione P, Galati S, Scarnati E, Peppe A, Stefani A. Implantation of human peduncolopontine nucleus: a safe and clinically relevant target in Parkinson's disease.

Neuroreport 2005; 16(17): 1877-81.

**IF. 1.805**

107) Filippi L, Manni C, Pierantozzi M, Brusa L, Danieli R, Stanzione P, Schillaci O. 123I-FP-CIT semi-quantitative SPECT detects preclinical bilateral dopaminergic deficit in early Parkinson's disease with unilateral symptoms.

Nucl Med Commun. 2006 May;26(5):421-6.

**IF. 1.315**

108) Stefani A, Galati S., Peppe A, Bassi A, Pierantozzi M, Hainsworth AH, Bernardi G, Orlacchio A, Stanzione P, Mazzone P. Spontaneous sleep modulates the firing pattern of parkinsonian subthalamic nucleus.

Exp Brain Res. 2006; 168(1-2): 277-80.

**IF. 2.256**

109) Brusa L, Petta F, Pisani A, Miano R, Stanzione P, Moschella V, Galati S, Finazzi Agrò E. Central acute D2 stimulation worsen bladder function in patients with mild Parkinson's disease.

J Urol. 2006; 175(1): 202-6; discussion 206-7.

**IF. 4.016**

110) Biroccio A, Del Boccio P, Panella M, Bernardini S, Di Ilio C, Gambi D, Stanzione P, Sacchetta P, Bernardini G, Martorana A, Federici G, Stefani A, Urbani A. Differential post-translational modifications of transthyretin in Alzheimer's disease: a study of the cerebro spinal fluid. *Proteomics*. 2006; 6(7): 2305-13. **IF. 4.426**

111) Marongiu R, Grezzi D, Ialongo T, Soleti F, Elia A, Cavone S, Albanese A, Altavista MC, Barone P, Brusa L, Cortelli P, Petrozzi L, Scaglione C, Stanzione P, Tinazzi M, Zeviani M, Dalla Piccola B, Bentivoglio AR, Valente EM, Garavaglia B, Italian PD Study Group. Frequency and phenotypes of LRRK2 mutation in Italian patients with Parkinson's disease. *Mov Dis* 2006; 21(8): 1232-5. **IF. 4.014**

112) Pierantozzi M, Pietroiusti A, Brusa L, Galati S, Stefani A, Lunardi G, Fedele E, Sancesario G, Bernardi G, Bergamaschi A, Magrini A, Stanzione P, Galante A. Helicobacter pilori eradication and l-dopa absorption in patients with PD and motor fluctuations. *Neurology* 2006; 66(12): 1824-29. **IF. 8.172**

113) Galati S, Mazzone P, Fedele E, Pisani A, Peppe A, Pierantozzi M, Brusa L, Tropepi D, Moschella V, Raiteri M, Stanzione P, Bernardi G, Stefani A. Biochemical and electrophysiological changes of substantia nigra pars reticulata driven by subthalamic stimulation in patients with Parkinson's disease. *Eur J Neurosci*. 2006; 23(11): 2923-28. **IF. 3.418**

114) Romigi A, Stanzione P, Marciani MG, Izzi F, Placidi F, Cervellino A, Giacomini P, Brusa L, Grossi K, Pierantozzi M. Effect of cabergoline added to levodopa treatment on sleep-wake cycle in idiopathic Parkinson's disease: an open label 24-hour polysomnographic study. *J Neural Transm*. 2006; 113(12): 1909-13. **IF. 2.259**

115) Stefani A, Fedele E, Galati S, Raiteri M, Pepicelli O, Brusa L, Pierantozzi M, Peppe A, Pisani A, Gattoni G, Hainsworth AH, Bernardi G, Stanzione P, Mazzone P. Deep brain stimulation in Parkinson's disease patients: biochemical evidence. *J Neural Transm Suppl*. 2006;(70):401-8. **IF. 2.525**

116) Brusa L, Versace V, Koch G, Iani C, Stanzione P, Bernardi G, Centonze D. Low frequency rTMS of the SMA transiently ameliorates peak-dose LID in Parkinson's disease. Clin Neurophysiol. 2006 Sep;117(9):1917-21.

**IF. 3.122**

117) Alimonti A, Ristori G, Giubilei F, Stazi MA, Pino A, Visconti A, Brescianini S, Monti MS, Forte G, Stanzione P, Bocca B, Bomboi G, D'Ippolito C, Annibali V, Salvetti M, Sancesario G. Serum chemical elements and oxidative status in Alzheimer's disease, Parkinson disease and multiple sclerosis. Neurotoxicology. 2007 May;28(3):450-6.

**IF. 2.918**

118) Stefani A, Lozano A, Peppe A, Stanzione P, Galati S, Tropepi D, Pierantozzi M, Brusa L, Scarnati E, Mazzone P. Bilateral deep brain stimulation of the pedunclopontine and subthalamic nuclei in severe Parkinson's disease.

Brain 2007; 130 (pt6) 1596-607

**IF. 9.490**

119) Brusa L, Petta F, Pisani A, Moschella V, Iani C, Stanzione P, Miano R, Finazzi-Agro E. Acute vs chronic effects of l-dopa on bladder function in patients with mild Parkinson disease. Neurology. 2007 May 1;68(18):1455-9.

**IF. 8.172**

120) Florio T, Scarnati E, Confalone G, Minchella D, Galati S, Stanzione P, Stefani A, Mazzone P. High-frequency stimulation of the subthalamic nucleus modulates the activity of pedunclopontine neurons through direct activation of excitatory fibres as well as through indirect activation of inhibitory pallidal fibres in the rat.

Eur J Neurosci. 2007 Feb;25(4):1174-86.

**IF. 3.418**

121) Centonze D, Koch G, Versace V, Mori F, Rossi S, Brusa L, Grossi K, Torelli F, Prosperetti C, Cervellino A, Marfia GA, Stanzione P, Marciani MG, Boffa L, Bernardi G. Repetitive transcranial magnetic stimulation of the motor cortex ameliorates spasticity in multiple sclerosis. Neurology. 2007 Mar 27;68(13):1045-50. .

**IF. 8.172**

122) Sallustio F, Diomedi M, Centonze D, Stanzione P.

Saving the ischemic penumbra: potential role for statins and phosphodiesterase inhibitors.

Curr Vasc Pharmacol. 2007 Oct;5(4):259-65. Review.

**IF. 2.970**

123) Mazzone P, Insola A, Lozano A, Galati S, Scarnati E, Peppe A, Stanzione P, Stefani A.  
Peripeduncular and pedunculo pontine nuclei: a dispute on a clinically relevant target.  
Neuroreport. 2007 Aug 27;18(13):1407-8. **IF. 1.805**

124) Romigi A, Placidi F, Peppe A, Pierantozzi M, Izzi F, Brusa L, Galati S, Moschella V, Marciani MG,  
Mazzone P, Stanzione P, Stefani A.  
Pedunculo pontine nucleus stimulation influences REM sleep in Parkinson's disease.  
Eur J Neurol. 2008 Jul;15(7):e64-5. **IF. 2.510**

125) Galati S, Scarnati E, Mazzone P, Stanzione P, Stefani A.  
Deep brain stimulation promotes excitation and inhibition in subthalamic nucleus in Parkinson's  
disease.  
Neuroreport. 2008;19(6):661-6. Err in: Neuroreport. 2008 ;19(8):899. **IF. 1.805**

126) Peppe A, Gasbarra A, Stefani A, Chiavalon C, Pierantozzi M, Fermi E, Stanzione P, Caltagirone  
C, Mazzone P.  
Deep brain stimulation of CM/PF of thalamus could be the new elective target for tremor in  
advanced Parkinson's Disease?  
Parkinsonism Relat Disord. 2008 Aug;14(6):501-4. **F. 2.4.6**

127) Pierantozzi M, Palmieri MG, Galati S, Stanzione P, Peppe A, Tropepi D, Brusa L, Pisani A,  
Moschella V, Marciani MG, Mazzone P, Stefani A.  
Pedunculo pontine nucleus deep brain stimulation changes spinal cord excitability in Parkinson's  
disease patients.  
J Neural Transm. 2008 May;115(5):731-5. **IF. 2.259**

128) Placidi F, Izzi F, Romigi A, Stanzione P, Marciani MG, Brusa L, Sperli F, Galati S, Pasqualetti P,  
Pierantozzi M.  
Sleep-wake cycle and effects of cabergoline monotherapy in de novo Parkinson's disease patients.  
An ambulatory polysomnographic study.  
J Neurol. 2008 Jul;255(7):1032-7. **IF. 2.930**

129) Antonini A, Colosimo C, Marconi R, Morgante L, Barone P; PRIAMO study group. The PRIAMO study: background, methods and recruitment.

Neurol Sci. 2008 Apr;29(2):61-5.

**IF. 1.120**

130) Sallustio F, Di Legge S, Koch G, Stanzione P.

Intracranial dissection and extracranial hypoplasia of the internal carotid artery.

J Ultrasound Med. 2008 May;27(5):795-8.

**IF. 1.181**

131) Galati S, D'angelo V, Scarnati E, Stanzione P, Martorana A, Procopio T, Sancesario G, Stefani A. In vivo electrophysiology of dopamine-denervated striatum: focus on the nitric oxide/cGMP signaling pathway.

Synapse. 2008 Jun;62(6):409-20.

**IF. 2.557**

132) Marongiu R, Ferraris A, Ialongo T, Michiorri S, Soleti F, Ferrari F, Elia AE, Ghezzi D, Albanese A, Altavista MC, Antonini A, Barone P, Brusa L, Cortelli P, Martinelli P, Pellicchia MT, Pezzoli G, Scaglione C, Stanzione P, Tinazzi M, Zecchinelli A, Zeviani M, Cassetta E, Garavaglia B, Dallapiccola B, Bentivoglio AR, Valente EM; Italian PD Study Group.

PINK1 heterozygous rare variants: prevalence, significance and phenotypic spectrum.

Hum Mutat. 2008 Apr;29(4):565.

**IF. 6.887**

133) Gelmetti V, Ferraris A, Brusa L, Romano F, Lombardi F, Barzaghi C, Stanzione P, Garavaglia B, Dallapiccola B, Valente EM.

Late onset sporadic Parkinson's disease caused by PINK1 mutations: clinical and functional study.

Mov Disord. 2008 Apr 30;23(6):881-5.

**IF. 4.014**

134) Diomedi M, Stanzione P, Sallustio F, Leone G, Renna A, Misaggi G, Fontana C, Pasqualetti P, Pietroiusti A.

Cytotoxin-associated Gene-A-positive Helicobacter pylori strains infection increases the risk of recurrent atherosclerotic stroke.

Helicobacter. 2008 Dec;13(6):525-31.

**IF. 2.851**

135) Lunardi G, Galati S, Tropepi D, Moschella V, Brusa L, Pierantozzi M, Stefani A, Rossi S, Fornai F, Fedele E, Stanzione P, Hainsworth AH, Pisani A.

Correlation between changes in CSF dopamine turnover and development of dyskinesia in Parkinson's disease.

Parkinsonism Relat Disord. 2009 Jun; 15(5): 383-89.

**IF. 2.406**

136) Stefani A, Peppe A, Pierantozzi M, Galati S, Moschella V, Stanzione P, Mazzone P.

Multi-target strategy for Parkinsonian patients: The role of deep brain stimulation in the centromedian-parafascicularis complex.

Brain Res Bull. 2009 Feb 16; 78 (2-3): 113-118.

**IF. 2.184**

137) Brusa L, Agrò EF, Petta F, Sciobica F, Torriero S, Lo Gerfo E, Iani C, Stanzione P, Koch G.

Effects of inhibitory rTMS on bladder function in Parkinson's disease patients.

Mov Disord. 2009 Feb 15;24(3):445-8.

**IF. 4.014**

138) Zanini S., Moschella V., Stefani A., Peppe A., Pierantozzi M., Galati S., Costa A., Mazzone P., Stanzione P.

Grammar improvement following deep brain stimulation of the subthalamic and the pedunculopontine nuclei in advanced Parkinson's disease: a pilot study.

Parkinsonism Relat Dis 2009 Sep; 15(8): 606-09.

**IF. 2.406**

139) Sallustio F, Di Legge S, Koch G, Ippoliti A, Mauriello A, Stanzione P.

Urgent carotid endarterectomy: the role of serial ultrasound studies in early detection of plaque rupture.

J Ultrasound Med. 2009 Feb;28(2):239-43.

**IF. 1.181**

140) Brusa L, Iani C, Ceravolo R, Galati S, Moschella V, Marzetti F, Stanzione P, Stefani A.

Implantation of the nucleus tegmenti pedunculo pontini in a PSP-P patient: Safe procedure, modest benefits.

Mov Disord. 2009 Oct 15; 24(13): 2020-22.

**IF. 4.014**

141) Galati S, Stanzione P, D'Angelo V, Fedele E, Marzetti F, Sancesario G, Procopio T, Stefani A.

The pharmacological blockade of medial forebrain bundle induces an acute pathological synchronization of the cortico-subthalamic nucleus-globus pallidus pathway.

J Physiol. 2009 Sep 15;587(Pt 18):4405-23. Epub 2009 Jul 21.

**IF. 4.764**

142) Koch G, Brusa L, Carrillo F, Lo Gerfo E, Torriero S, Oliveri M, Mir P, Caltagirone C, Stanzione P. Cerebellar magnetic stimulation decreases levodopa-induced dyskinesias in Parkinson disease.

Neurology. 2009 Jul 14;73(2):113-9.

**IF. 8.172**

143) Lunardi G., Galati S., Tropepi D., Morchella V., Brusa L., Pierantozzi M., Stefani A., Rossi S., Fornai F., Fedele E., Stanzione P., Hainsworth AH., Pisani A.

Correlation between changes in CSF dopamine turnover and development of dyskinesia in Parkinson's disease.

Parkinsonism Relat Disord 2009; 15(5): 383-89.

**IF. 2.406**

144) Sallustio F., Rotondo F., Di Legge S., Stanzione P.

Cilostazol in the management of atherosclerosis.

Curr Vasc Pharmacol 2010; 8(3): 363-72.

**IF. 2.970**

145) Sallustio F., Di Legge S., Rossi C., Stanzione P.,

Ultrasound evaluation of early changes in arterial dissection.

J Stroke Cerebrovasc. Dis. 2010; 19(2): 167-68.

146) Pisani V., Moschella V., Bari M., Fezza F., Galati S., Bernardi G., Stanzione P., Pisani A., Maccarrone M.

Dynamic changes of anandamide in the cerebrospinal fluid of Parkinson's disease patients.

Mov Dis. 2010; 25(7): 920-24.

**IF. 4.014**

147) Stefani A., Pierantozzi M., Ceravolo R., Brusa L., Galati S., Stanzione P.

Deep brain stimulation of peduncolopontine nucleus (PPTg) promotes cognitive and metabolic changes: a target-specific effect or response to a low frequency pattern of stimulation ?

Clin EEG Neurosci. 2010; 41(2): 82-86.

**IF. 1.121**

148) Sallustio F., Di Legge S., Rizzato B., Stanzione P., Koch G.

Changes in cerebrovascular reactivity following low-frequency repetitive transcranial magnetic stimulation.

J Neurol Sci 2010; 295(1-2): 58-61.

**IF. 2.342**

149) Galati S., D'Angelo V., Olivola E., marzetti F., Di giovanni G., Stanzione P., Stefani A.

Acute inactivation of medialforbrain bundle imposes oscillations in the SNr: a challenge for 6-OHDA model ?

Exp Neurol. 2010;225(2): 294-301.

**IF. 3.914**

150) Assogna F., Pontieri FE., Cravello L., Peppe A., Pierantozzi M., Stefani A., Stanzione P., Pellicano C., Caltagirone C., Spalletta G.

J Int Neuropsychol Soc. 2010; 16(5). 867-76.

**IF. 2.766**

151) Peppe A., Pierantozzi M., Chiavalon C., Marchetti F., Caltagirone C., Musico M., Stanzione P., Stefani A.

Deep braij stimulation of peduncolopontine tegmentum and subthalamic nucleus: effects on gait in Parkinson's disease.

Gait Posture 2010; 32(4): 512-18.

**IF. 2.576**

152) Sallustio F., Di Legge S., Marziali S., Ippoliti A., Stanzione P.

Floating carotid thrombus treated by intravenous heparin and endoarteriectomy.

J Vasc Surg. 2010; Oct 8 e pub ahead

**IF. 3.517**

153) Pisani V., Stanzione P., Meschini A., Marziali S., Garaci FG., Brancati F., Nicolatti A., Zappia M., Pisani A.

Putaminal but not nigral alterations characterize emiparkinsonism-hemiatrophy syndrome: a case report.

Mov Dis 2010: Oct 13 epub ahead

**IF. 4.014**

154) Ceravolo R., Brusa L., Galati S., Volterrani D., Peppe A., Siciliano G., Pierantozzi M., Morchella V., Bonuccelli U., Stanzione P., Stefani A.



Low frequency stimulation of the nucleus tegmenti peduncolopontini increases cortical metabolism in Parkinsonian patients.

Eur J Neurol 2010 Nov 19 epub ahead

**IF. 2.510**

155) Pisani V., Madeo G., Tassone A., Sciamanna G., Maccarrone M., Stanzione P., Pisani A.  
Homeostatic changes of the endocannabinoid system in Parkinson's disease

Mov Dis. 2010 Dec 13 epub ahead.

**IF. 4.014**

**Tot 272.106**

**Paper da 71 a 155 n= 86 meno 2 libri 84**

**Mean 3.23**

**Dal 2005 al 2010**

**Papers da 96 a 154**

**Tot 232,463**

**Mean 4.0079**